



PRIDE SEEDS

SILAGE CORN

A4477HM

2150-2250 CHU



Product Features

- Conventional silage, high-moisture and grazing corn, offering higher grain density and disease tolerance
- Features slow grain-drying rate and preserves reliable and consistent feed quality at ideal moisture content
- Aggressive spring vigour for early maturity zone
- High starch levels provide excellent quality silage
- Excellent for grazing use with high yield, nutrition and strong stalks
- Produces very girthy, exceptionally consistent ears, producing flint kernels on white cob
- A solid fit for beef or dairy operations

Agronomic Characteristics

| | |
|--------------------|------|
| EDF or EDP | EDF |
| Flowering | A |
| Mid Flowering GDU | 1120 |
| Plant Height | M |
| Emergence | G |
| Spring Vigour | G |
| Stalk Strength | EXC |
| Root Strength | EXC |
| Staygreen | EXC |
| Drought Tolerance | VG |
| Plant Health | EXC |
| Ear Type | FL |
| Starch | VG |
| Crude Protein | EXC |
| NDFD | EXC |
| Milk or Beef Acre | VG |
| Milk or Beef Tonne | EXC |
| Goss's Wilt | VG |

Ratings EXC = Excellent VG = Very Good G = Good F = Fair

Plant Height S = Short M = Medium T = Tall V/T = Very Tall

Flowering E = Early E/A = Early Average A = Average L = Late

Ear Type F = Fixed SF = Semi-Flex FL = Flex



PRIDE SEEDS

WWW.PRIDSEEDS.COM

All orders and sales are subject to the PRIDE Seeds Terms and Conditions of Sale, which include but are not limited to the Limitation of Warranty & Remedy and Agronomic Zone and Planting Year. All Terms and Conditions of Sale are subject to change from time to time without prior notice. PRIDE® & Design, P® & Design, Advantage Acre®, and Pride Advantage Acre® & Design are registered trademarks of AgReliant Genetics Inc. Characteristics are assigned by PRIDE® based on comparisons with other PRIDE® products (not competitive products) through in-house field testing. Individual results may vary, and performance may vary from location to location and from year to year. This result may not be an indicator of results you may obtain as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible.